

Using the iPhone for Voice Recording in Laryngology

Jeremy Hornibrook,¹ Emily Lin,² & Tika Ormond²


¹Department of Otolaryngology, Christchurch Hospital

*²Department of Communication Disorders, University of Canterbury
Christchurch, New Zealand*




Background

- iPhone (first released in: Year 2007): a multi-media enabled mobile phone with advanced
 - Computing capabilities
 - Connectivity: Internet and email access
- iPhone 3G & up: sampling rate = 48,000 Hz (Lossless)





Research Question

- Can the iPhone be used to record voices for acoustic analysis to:
 - Identify voice aberrations
 - Monitor voice changes: e.g.,
pre- and post-treatment differences?
- 

Participants


- Twenty-two voice patients (10 males & 12 females), aged 25-92 years (Mean = 54.8, SD = 18.5), including:
 - 10 patients (6 males & 4 females) who underwent phonosurgery:
 - Aged 33-79 years (Mean = 47.6, SD = 15.3)
 - Pathology:
 - **Mass Lesions** (8 cases): cyst (2), nodules (2), edema (1), papilloma (1), polyps (1), benign tumor (1)
 - Treatment: microsurgery
 - **Vocal Paralysis** (2 cases)
 - Treatment: medialization laryngoplasty

Participant's Task

- To read the first 6 sentences of “Rainbow passage” (*Fairbanks, 1960*)
- Recordings:
 - Before treatment: for all of the 22 participants
 - After treatment: for 10 of the participants
 - Time between pre- and post-surgery recordings:
22-259 days (Mean = 103, SD = 95.5)


Instrumentation

- Apple iPhone (Model A1303)
- Wav MP3 converter software (Hoo Technologies, USA)
- TF32 voice analysis software (*Milenkovic, 1987*)



Procedure

- Laryngostroboscopic examination
- Acoustic recording



Case One: Left Vocal Fold Paralysis

- Male, aged 79 years, aortic arch aneurysm




 PRE


 POST (medialization laryngoplasty)

Case Two: Polyp on LVF

- Male, aged 33 years




 PRE




 POST (microsurgery)

Case Three: Inflammatory myoblastic tumor

- Male, aged 34 years




 PRE




 POST (microsurgery)

Case Four: Mass lesion on RVF

- Male, aged 46 years




 PRE



 POST (microsurgery)

Case Five: Nodules

- Female, aged 39 years



PRE



POST (microsurgery)


Data Analysis



Vowel-based measures: F0, perturbation measures (%Jit, %Shim, SNR), H1-H1, SPR, F1, F2

Sentence-based measures: ST

Data Analysis



Vowel-based measures: F0, perturbation measures (%Jit, %Shim, SNR), H1-H1, SPR, F1, F2

Sentence-based measures: ST

Measurement

- **Vowel-based measures** (time waveform analysis):
 - Fundamental frequency (**F0**):
 - Edema: decreased F0 (*Sorensen & Horii, 1982*)
 - Speaking F0 changes after treatment of functional voice (*Roy & Taskco, 1994*)

Changes in mass and stiffness → Change in F0
 - Perturbation measures:
 - Jitter (or percent jitter; **%Jit**): cycle-to-cycle pitch variation
(e.g., *Lieberman, 1961; Eskenazi et al., 1990; Dejonckere et al., 1996; Wolfe & Martin, 1997; Bhuta et al., 2004*)
 - Shimmer (or percent shimmer, **%Shim**): cycle-to-cycle amplitude variation
(e.g., *Dejonckere et al., 1996; Wolfe & Martin, 1997; Bhuta et al., 2004*)
 - Signal-to-noise ratio (**SNR**): energy ratio between periodic & aperiodic components
(*Yanagihara, 1967; Wolfe & Martin, 1997; Brockmann, Storck, Carding, & Drinnan, 2008*)

Less hoarse → decreased jitter & shimmer; increased SNR



Measurement – continued

- **Vowel-based measures** (spectral analysis):
 - Amplitude difference between the first two harmonics (**H1-H2**) as measured from a spectrum (without pre-emphasis)

Less breathy voice → Smaller H1-H2 (i.e. less H1 dominance)

(Klatt & Klatt, 1990; Hillenbrand, Cleveland, & Erickson, 1994; de Krom, 1995; Hillenbrand & Houde, 1996; Stone, Cleveland, Sundberg, & Prokop, 2003)

Amplitude



PRE (M79-paralysis)




POST-surgery

Measurement – continued

- **Vowel-based measures** (spectral analysis) - *continued*:
 - Singing power ratio (**SPR**):
 - Defined as: amplitude difference between the highest spectral peak between 0 and 2 kHz and that between 2 and 4 kHz as measured from a spectrum (with pre-emphasis)

Greater voice projection power → Smaller SPR

(Omori, Kacker, Carroll, Riley, & Blaugrund, 1996)



PRE (M46-paralysis)




POST-surgery

Measurement – continued

- **Vowel-based measures** (spectral analysis) - *continued*:
 - Formants One and Two frequencies (**F1** and **F2**):
 - Defined as: the highest two spectral peaks in a LPC (linear predictive coding) spectrum
 - Related to vocal tract configuration (constriction or tongue placement)
 - Space enclosed by the corner vowels /i, a, u/ in a F1-F2 plot = **vowel space**

Increased vowel differentiation → Larger vowel space area

(Bradlow, Toretta, & Pisoni, 1996; Roy, Nissen, Dromey, & Sapis, 2009; Turner, Tjaden, & Weismer, 1995; Weismer, Jeng, Laures, Kent, & Kent, 2001)



• PRE (M34-tumor)



/i/



/a/



/u/

• POST-surgery



/i/



/a/



/u/


Measurement – continued

- **Sentence-based measures:**

- Spectral tilt (**ST**): amplitude difference between the highest spectral peak between 0 and 1 kHz and that between 1 and 5 kHz as measured from a LTA (long-time average) spectrum (without pre-emphasis)

More lax vocal fold adduction → higher ST (i.e., steeper slope)

(Löfqvist, 1987; Mendoza et al., 1996)



PRE (F67-edema)



POST-surgery

Reliability

- Automatic computer derivations of acoustic measures: **100% reliability**, except for errors due to variation in vowel segmentation.
- **One third** of the data for the ten patients with pre- and post-treatment recordings were re-segmented and analyzed. The measure-remmeasure reliabilities (Pearson's r) were **high** for all vowel-based measures:


Measure	n	r
F0	20	0.99
%Jit	20	0.99
%Shim	20	0.98
SNR	20	0.97
F1	20	0.93
F2	20	0.88
H1-H2	20	0.97
SPR	20	0.96

Statistical Analysis

- A series of two-way (2 groups X 3 vowels) mixed model ANOVAs on F0, %Jit, %Shim, SNR, H1-H2, SPR, F1, & F2.
 - Between-subject variable: group (Pre vs. Post)
 - Within-subject variable: vowel (/i/, /a/, & /u/)
- A series of paired t tests on vowel space area & ST (averaged from six sentences).
- Significance level set at 0.05.


Results

- After surgery: Significantly lower F0, %Jit, & %Shim and higher SNR & vowel space area.
 - Vowel space area: significant larger vowel space area ($t = -3.746$, $df = 9$, $p = 0.007$)
 - ANOVA results:
 - No significant vowel by group interaction effect.
 - Significant vowel effect on F0, SNR, SPR, F1, & F2
 - Significant group effect on F0, %Jit, %Shim, & SNR.
 - Shown below with the average pre- and post-surgery scores normalized based on the pre-treatment data of 22 patients:



Results - continued

- F0 & H1-H2 change varied by pathology:
 - Mass lesion (8 cases): After surgery,
 - Lower pitch (except for edema): Significantly lower F0 as a whole [$F(1, 14) = 6.788, p = 0.035, \eta^2 = 0.08$].
 - No significant H1-H2 change
 - Paralysis (2 cases): After surgery,
 - No significant F0 change
 - Less breathy: Significantly lower H1-H2 [$F(1, 2) = 996.755, p = 0.002, \eta^2 = 0.71$]



Results - *continued*

- **SPR change varied by vowel:**
 - With the vowel /a/, the majority (7 out of 10) showed a high preoperative SPR value (i.e., less voice projection power)
 - **More voice projection power:** This subgroup showed a significantly lower SPR after surgery ($t = 3.383$, $df = 6$, $p = 0.015$).
- **ST change varied by gender:** After surgery,
 - Females ($n = 4$):
 - **More lax vocal fold adduction:** Significantly higher ($t = -7.683$, $df = 3$, $p = 0.005$).
 - Males ($n = 6$): no significant change

Discussion - Continued

- Signs of voice improvement:
 - The decrease in %Jit & %Shim and the increase in SNR found in the postoperative voices were expected as previous studies have shown that **phonatory stability** could be compromised by vocal pathology and improved with effective treatment.
 - The expansion of vowel space area after surgery reflects improved **speech clarity**, suggesting that voice quality may affect vowel intelligibility.
- **No consistent F0 change** after surgery for patients with vocal fold paralysis: this agrees with previous findings (LaBlance & Maves, 1992).

Limitations

- Small sample size
- Observations made in the subgroups regarding the confounding effects of **pathology**, **vowel**, and **gender** on the pre- and post-surgery acoustic changes require further investigations

Conclusions

- Voice recordings using iPhone are **adequate** for voice recording in acoustic assessment of voice quality in .
- However, due to large inter-subject variations, most of these measures are more useful for **intra-subject comparison** (to monitor changes within individuals) than for norm-referenced comparisons.

References

- Bhuta, T., Patrick, L., & Garnett, J. D. (2004). Perceptual evaluation of voice quality and its correlation with acoustic measurements. *Journal of Voice*, 18, 299-304.
- Bradlow, A. R., Torretta, G. M., and Pisoni, D. B. (1996). Intelligibility of normal speech I: global and fine grained acoustic-phonetic talker characteristics. *Speech Communication*, 20, 255-272.
- Brockmann, M., Storck, C., Carding, P. N., & Drinnan, M. J. (2008). Voice loudness and gender effect on jitter and shimmer in healthy adults. *Journal of Speech, Language, and Hearing Research*, 51, 1152-1160.
- de Krom, G. (1995). Some spectral correlates of pathological breathy and rough voice quality for different types of vowel fragments. *Journal of Speech and Hearing Research*, 38, 794-811.
- Dejonckere, P. H., Remacle, M., Fresnel-Ebaz, E., Woisard, V., Crevier-Buchman, L., & Millet, B. (1996). Differentiated perceptual evaluation of pathological voice quality: reliability and correlations with acoustic measurements. *Rev Laryngol Otol Rhinol (Bord)*, 117, 219-224.
- Eskenazi, L., Childers, D. G., & Hicks, D. (1990). Acoustic correlates of vocal quality. *Journal of Speech and Hearing Research*, 33, 298-306.
- Fairbanks, G. (1960). *Voice and Articulation Drillbook*. 2nd ed. New York: Harper & Row. Pg. 127.

References - continued

- Hillenbrand, J., Cleveland, R. A., & Erickson, R. L. (1994). Acoustic correlates of breathy voice quality. *Journal of Speech and Hearing Research*, 37, 769-778.
- Hillenbrand, J., & Houde, R. A. (1996). Acoustic correlates of breathy vocal quality: dysphonic voices and continuous speech. *Journal of Speech and Hearing Research*, 39, 311-321.
- Klatt, D. H., & Klatt, L. C. (1990). Analysis, synthesis and perception of voice quality variations among female and male talkers. *Journal of the Acoustical Society of America*, 87, 820-854.
- Kotby, M. N., Titze, I. R., Saleh, M. M., & Berry, D. A. (1993). Fundamental frequency stability in functional dysphonia. *Acta Otolaryngologica*, 113, 4439-444.
- LaBlance G. R., & Maves, M. D. (1992). Acoustic characteristics of post-thyroplasty patients. *Otolaryngol Head Neck Surg*, 107, 558-563.
- Löfqvist, A., & Mandersson, B. (1987). Long-time average spectrum of speech and voice analysis. *Folia Phoniatica (Basel)*, 21, 221-229.
- Mendoza, E., Munoz, J., Valencia Naranjo, N. (1996). The long-term average spectrum as a measure of voice stability. *Folia Phoniatica et Logopaedica*, 48, 57-64.

References - continued

- Omori, K., Kacker, A., Carroll, L. M., Reiley, W. D., & Blaugrund, S. M. (1996). Singing power ratio: quantitative evaluation of singing voice quality. *Journal of Voice*, 10, 228-235.
- Roy, N., Nissen, S. L., Dromey, C., & Sapis, S. (2009). Articulatory changes in muscle tension dysphonia: evidence of vowel space expansion following manual circumlaryngeal therapy. *Journal of Communication Disorders*, 42, 124-135.
- Roy, N., & Tasko, S. M. (1994). Speaking fundamental frequency (SFF) changes following successful management of functional dysphonia. *Journal of Speech-Language Pathology and Audiology*, 18, 115-120.
- Sorensen, D., & Horii, Y. (1982). Cigarette smoking and voice fundamental frequency. *Journal of Communication Disorders*, 15, 135-144.
- Stone, R. E. Jr., Cleveland, T. F., Sundberg, P. J., & Prokop, J. (2003). Aerodynamic and acoustical measures of speech, operative, and Broadway vocal styles in a professional female singer. *Journal of Voice*, 17, 283-297.

References - continued

- Turner, G. S., Tjaden, K., & Weismer, G. (1995). The influence of speaking rate on vowel space and speech intelligibility for individuals with amyotrophic lateral sclerosis. *Journal of Speech and Hearing Research*, 38, 1001-1013.
- Watts, C. Barnes-Burroughs, K., Estis, J., Blanton, D. (2006). The singing power ratio as an objective measure of singing voice quality in untrained talented and nontalented singers. *J. of Voice*, 20, 82-88.
- Weismer, G., Jeng, J-Y., Laures, J. S., Kent, R. D., & Kent, J. F. (2001). Acoustic and intelligibility characteristics of sentence production in neurogenic speech disorders. *Folia Phoniatrica*, 53, 1-18.
- Winholtz, W. S., & Titze, I. R. (1998). Suitability of minidisc (MD) recordings for voice perturbation analysis. *J. of Voice*, 12, 138-142.
- Wolfe, V. & Martin, D. (1997). Acoustic correlates of dysphonia: type and severity. *Journal of Communication Disorders*, 30, 403-416.
- Yanagihara, N. (1967). Significance of harmonic changes and noise components in hoarseness. *J Speech Hear Res*, 10, 531-541.